Developmental changes in NMDA-induced cell swelling and its transition to necrosis measured with ^{1}H magnetic resonance imaging, impedance and histology (93) 109
Van Maele-Fabry, G., see Gofflot, F. (93) 62
Verheul, H.B., see Van Lookeren Campagne, M. (93) 109
Vermeulen, J.P., see Van Lookeren Campagne, M. (93) 109
Wadhwa, S., see Nag, T.C. (93) 23
Walkley, S.U., see Goodman, L.A. (93) 162
Wang, X.M., see Terman, J.R. (93) 33
Waters, N.S., see Denenberg, V.H. (93) 100
Yager, J.Y., Shuaib, A. and Thornhill, J. The effect of age on susceptibility to brain damage in a model of global hemispheric hypoxia-ischemia (93) 143
Yamamoto, M., McCaffery, P. and Dräger, U.C. Influence of the choroid plexus on cerebellar development: analysis of retinoic acid synthesis (93) 182
Zupanc, G.K.H., see Stroh, T. (93) 76

(contents continued)

Short communications

Developmental expression of Fos-lacZ in the brains of postnatal transgenic rats
G.M. Kasof, R.J. Smeyne, T. Curran and J.I. Morgan (USA) 191
All-trans retinoic acid affects the expression of orphan receptors COUP-TF I and COUP-TF II in the developing neural tube
K. Brubaker, M. McMillan, T. Neuman and H.O. Nornes (USA) 198

Author Index 203